

# Draft Environment and Social Impact Assessment

Project Number: 55205-001  
29 April 2022

## Lao PDR: Monsoon Wind Power Project Part 20: Appendix F (part 6)

Prepared by Impact Energy Asia Development Limited (IEAD) for the Asian Development Bank.

This draft environment and social impact assessment is a document of the borrower. The views expressed herein do not necessarily represent those of ADB's Board of Directors, Management, or staff, and may be preliminary in nature. Your attention is directed to the [“terms of use”](#) section of ADB's website.

In preparing any country program or strategy, financing any project, or by making any designation of or reference to a particular territory or geographic area in this document, the Asian Development Bank does not intend to make any judgments as to the legal or other status of any territory or area.



# Monsoon Wind Power Project, Sekong and Attapeu Provinces, Lao PDR

Environmental and Social Impact  
Assessment

29 April 2022

Project No.: 0598121

Document details	
Document title	Monsoon Wind Power Project, Sekong and Attapeu Provinces, Lao PDR
Document subtitle	Environmental and Social Impact Assessment
Project No.	0598121
Date	29 April 2022
Version	2.0
Author	Aurora Finiguerra, Cheryl Ng, Elaine Wong, Hoa Tran, Jacopo Ventura, Mingkwan Naewjampa, Shubhankar Khare, Tirapon Premchitt, Winee Tammaruk
Client Name	Impact Energy Asia Development Limited (IEAD)

#### Document history

Version	Revision	Author	Reviewed by	ERM approval to issue		Comments
				Name	Date	
1	1.1	As above	Kamonthip Ma-Oon, Sabrina Genter, Les Hatton, George Chatziannidis, Simone Poli, Aniket Jalgaonkar	Kamonthip Ma-Oon	18-02-22	Draft to IEAD
1	1.2	As above	As above	Kamonthip Ma-Oon	25-02-22	Draft to IEAD
1	1.3	As above	As above	Kamonthip Ma-Oon	23-03-22	Draft to IEAD and ADB
1	1.4	As above	As above	Kamonthip Ma-Oon	30-03-22	Draft to IEAD and ADB
1	1.4	As above	As above	Kamonthip Ma-Oon	21-04-22	Draft to IEAD and ADB
2	2.1	As above	As above	Kamonthip Ma-Oon	29-04-22	Final ESIA Report

## **APPENDIX F      BIODIVERSITY BASELINE SURVEY REPORTS**



### Annex 3b. List of herpetofauna (amphibian) records from the survey area

#	Common Name	Scientific Name	Local Name	Recorded by	IUCN Redlist	Survey block					Survey season		Abundance	Comments
						SB1	SB2	SB3	SB4	SB45	Wet	Dry		
		<b>Bufonidae</b>												
1		<i>Duttaphrynus melanostictus</i>			LC					1	x			Very common in open/ disturbed area. Dakcheung to
2		<i>Ingerophrynus galeatus</i>			LC			1	2		x		2	
		<b>Ranidae</b>												
3		<i>Odorrana morafkai</i>			LC	5	5	5	5	2	x		5	
4		<i>Odorrana banaorum</i>			LC	3	3	3	4		x		4	
5		<i>Odorrana khalam</i>			LC	1					x		1	
6		<i>Odorrana absita</i>			LC			1			x		1	
7		<i>Odorrana</i> sp.						1			x		1	With green on limbs
8		<i>Papurana milleti</i>			LC		2	3	5	2	x		4	
9		<i>Papurana attigua</i>			LC	3		3	2	1	x		3	
10		<i>Sylvirana montosa</i>			LC				1		x	x		
11		<i>Amolops spinaepectoralis</i>			LC			3			x		2	First record of Laos! Found only locally.
		<b>Dicoglossidae</b>												
12		<i>Fejervarya limnocharis</i>			LC					1	x	x	4	Abundant in disturbed areas.
13		<i>Quasipaa</i> sp.									x			Undescribed species
14		<i>Limnonectes cf taylori</i>			LC		1				x	x		
15		<i>Limnonectes kizirianii</i>			LC		1				x	x		
16		<i>Limnonectes poilani</i>			LC						x			First record of Laos!
17		Spiny torrent						3	4	2		x	4	
		<b>Megophryidae</b>												
18		<i>Ophryophryne hansi</i>			DD	4	3	4	5	1	x		4	Common along forest streams
19		<i>Ophryophryne poilani</i>						1			x			
20		<i>Leptobranchella firsi</i>				2					x			First record of Laos!
21		<i>Leptobranchella</i> unid.									x			
22		<i>Brachytarsophrys intermedia</i>	Found by local people		LC		2				x			Near Dakchung township
23		<i>Xenophrys cf maosonensis</i>					3		5		x		4	Undescribed species
		<b>Microhylidae</b>												
24		<i>Microhyla berdmorei</i>									x			
25		<i>Microhyla butleri</i>									x			Only found in open disturbed area
26		<i>Microhyla heymonsi</i>									x			Only found in open disturbed area
27		<i>Micryletta</i> sp.						1			x			Only found in open disturbed area
28		<i>Nanohyla marmorata</i>			LC	1		2	4	1	x			
29		<i>Nanohyla nanapollexa</i>		Just observed	DD	1					x			

		<b>Rhacophoridae</b>											
30		Raorchestes sp.					1				x		
31		Kurixalus ballogaster	LC	1		1	3	2			x		More common at open areas
32		Kurixalus odontotarsus	LC				5	1			x		Locally distributed
33	Serrate Frilled Treefrog	Kurixalus cf gryllus	VU				1				x	1	Seems rare
34		Zhangixalus feae	LC	1		1					x	2	
35		Rhacophorus robertingeri	LC	3		3			1		x	4	
36		Rhacophorus sp nov				1					x		Undescribed species. Found in open area
37		Theioderma gordonii	LC					1			x		
38		Theioderma truongsoneensis	DD					1			x		First record of Laos!
39		Gracixalus supercornutus	NT	1				2			x		Locally fairly abundant e.g northern Phou Koungking
40		Polypedates megachephalus									x		Common species of open and disturbed habitats.
41		Polypedates impresus				1	1				x		
42		Polypedates sp.?				1					x		Unusual red coloration, appears overall different
Remarks: Abundance: 1. Rare/Low = 1; 2. Occasional = 2-5; 3. Frequent =5-10; 4. Common =10-20; 5. Abundant >20													

**Annex 4. Photos of some other wildlife species from the survey area**



4a. *Dopasia sokolovi* (second record of Laos) from SB3



4b. Feeding site of Pangolins (Chinese Pangolin)



4c. Claw marks of Bears on tree trunk (Sun Bear) from SB5



4d. Bat sp. from SB3



4e. *Gymnure* sp. from SB5



4f. Sokolov's Grass Lizard from SB3 (second record of Laos)

Annex 5. Photos of some forest landscape and structures from the survey area



Survey block #1 (Southern Annamite)



Survey block #2 (Southern Annamite)



Survey block #3 (Phou Koungking East)



Survey block #4 (Phou Koungking West)



Survey block #5 (Phou Yai)

## Annex 6. Waypoints for important (key and GT species) records

1 Plants: Coordinates of first and second records, Globally Threatened and New Species Candidates										
2	3 Group and Family	Species	GT	Endemism Species			Survey block	UTM		Alt (m a.s.l.)
				First Record	Second Record	New SP Candidat		X	Y	
4	Pentaphragaceae	Adinandra integerrima T.Anderson ex Dyer		x			4/2	728974	1712562	
5	Pentaphragaceae	Adinandra integerrima T.Anderson ex Dyer		x			4/3	729218	1712797	
6	Pentaphragaceae	Adinandra integerrima T.Anderson ex Dyer		x			4/4	730017	1712249	
7	Pentaphragaceae	Adinandra integerrima T.Anderson ex Dyer		x			4/5	730445	1712019	
8	Pentaphragaceae	Adinandra integerrima T.Anderson ex Dyer		x			4/6	731205	1712548	
9	Lauraceae	Alseodaphne bidouensis Yahara		x			2/1	750740	1722688	
10	Lauraceae	Alseodaphne bidouensis Yahara		x			2/3	751025	1722672	
11	Marattiaceae	Angiopteris wangii Ching		x			2/6	752943	1720599	
12	Icacinaceae	Apodytes dimidiata E.Mey. ex Arn.		x			3/1	734837	1712290	
13	Icacinaceae	Apodytes dimidiata E.Mey. ex Arn.		x			3/2	735209	1712032	
14	Icacinaceae	Apodytes dimidiata E.Mey. ex Arn.		x			3/3	735792	1712515	
15	Primulaceae	Ardisia gracilenta C.M.Hu & J.E.Vidal		x			3/2	735209	1712032	
16	Primulaceae	Ardisia gracilenta C.M.Hu & J.E.Vidal		x			3/3	735792	1712515	
17	Primulaceae	Ardisia gracilenta C.M.Hu & J.E.Vidal		x			3/4	731614	1712736	
18	Primulaceae	Ardisia gracilenta C.M.Hu & J.E.Vidal		x			4/2	728974	1712562	
19	Primulaceae	Ardisia gracilenta C.M.Hu & J.E.Vidal		x			4/4	730017	1712249	
20	Primulaceae	Ardisia gracilenta C.M.Hu & J.E.Vidal		x			5/3	722916	1711735	
21	Primulaceae	Ardisia gracilenta C.M.Hu & J.E.Vidal		x			5/5	722080	1713133	
22	Rubiaceae	Argostemma bariense Pierre ex Pit.		x			2/6	752943	1720599	
23	Rubiaceae	Brachytome wallichii Hookf.		x			4/1	728886	1712825	
24	Rubiaceae	Brachytome wallichii Hookf.		x			4/2	728974	1712562	
25	Rubiaceae	Brachytome wallichii Hookf.		x			4/4	730017	1712249	
26	Melastomataceae	Bredia sp. (new genus record for Laos)		x			2/5	723254	1710986	
27	Theaceae	Camellia sp. 1				x	5/1	723071	1710374	
28	Theaceae	Camellia sp. 1				x	5/6	722534	1713764	
29	Rubiaceae	Chassalia curviflora (Wall.) Thwaites		x			3/1	734837	1712290	
30	Daphniphyllaceae	Daphniphyllum beddomei Craib		x			3/1	734837	1712290	
31	Daphniphyllaceae	Daphniphyllum beddomei Craib		x			3/2	735209	1712032	
32	Daphniphyllaceae	Daphniphyllum beddomei Craib		x			3/3	735792	1712515	
33	Daphniphyllaceae	Daphniphyllum beddomei Craib		x			3/4	731614	1712736	
34	Daphniphyllaceae	Daphniphyllum beddomei Craib		x			3/5	731869	1712562	
35	Daphniphyllaceae	Daphniphyllum beddomei Craib		x			3/6	732041	1712500	
36	Elaeocarpaceae	Elaeocarpus dubius DC.		x			1/3	752318	1717523	
37	Elaeocarpaceae	Elaeocarpus dubius DC.		x			1/4	751395	1717920	
38	Elaeocarpaceae	Elaeocarpus dubius DC.		x			2/1	750740	1722688	
39	Elaeocarpaceae	Elaeocarpus dubius DC.		x			2/2	750569	1723483	
40	Elaeocarpaceae	Elaeocarpus dubius DC.		x			2/4	752726	1722359	
41	Elaeocarpaceae	Elaeocarpus limitaneus Hand.-Mazz.		x			2/5	752836	1721953	
42	Gnetaceae	Gnetum gnemon L.		x			3/1	734837	1712290	
43	Euphorbiaceae	Gymnanthes remota (Steenis) Esser		x			4/1	728886	1712825	
44	Euphorbiaceae	Gymnanthes remota (Steenis) Esser		x			4/2	728974	1712562	
45	Euphorbiaceae	Gymnanthes remota (Steenis) Esser		x			4/3	729218	1712797	
46	Euphorbiaceae	Gymnanthes remota (Steenis) Esser		x			4/4	730017	1712249	
47	Euphorbiaceae	Gymnanthes remota (Steenis) Esser		x			4/5	730445	1712019	
48	Euphorbiaceae	Gymnanthes remota (Steenis) Esser		x			4/6	731205	1712548	
49	Araliaceae	Heptapleurum cambodianum (Yahara & Tagane) Lowry & G.M		x			1/4	751395	1717920	
50	Araliaceae	Heptapleurum cambodianum (Yahara & Tagane) Lowry & G.M		x			1/6	752607	1719693	
51	Araliaceae	Heptapleurum cambodianum (Yahara & Tagane) Lowry & G.M		x			3/4	731614	1712736	
52	Araliaceae	Heptapleurum cambodianum (Yahara & Tagane) Lowry & G.M		x			3/5	731869	1712562	
53	Araliaceae	Heptapleurum cambodianum (Yahara & Tagane) Lowry & G.M		x			5/3	722916	1711735	
54	Araliaceae	Heptapleurum cambodianum (Yahara & Tagane) Lowry & G.M		x			5/5	722080	1713133	
55	Araliaceae	Heptapleurum cambodianum (Yahara & Tagane) Lowry & G.M		x			5/6	722534	1713764	
56	Aquifoliaceae	Ilex chapaensis Merr.		x			4/6	731205	1712548	
57	Aquifoliaceae	Ilex chapaensis Merr.		x			5/1	723071	1710374	

58	Aquifoliaceae	Ilex chapaensis Merr.		x		5/3	723254	1710986
59	Aquifoliaceae	Ilex chapaensis Merr.		x		5/3	722916	1711735
60	Lauraceae	Lindera bokorensis Yahara & Tagane		x		1/5	751734	1717636
61	Lauraceae	Lindera bokorensis Yahara & Tagane		x		2/6	752943	1720599
62	Lauraceae	Lindera bokorensis Yahara & Tagane		x		4/1	728886	1712825
63	Lauraceae	Lindera bokorensis Yahara & Tagane		x		4/3	729218	1712797
64	Lauraceae	Lindera bokorensis Yahara & Tagane		x		4/4	730017	1712249
65	Lauraceae	Lindera bokorensis Yahara & Tagane		x		4/5	730445	1712019
66	Fagaceae	Lithocarpus elephantum (Hance) A.Camus		x		3/2	735209	1712032
67	Fagaceae	Lithocarpus elephantum (Hance) A.Camus		x		3/5	731869	1712562
68	Fagaceae	Lithocarpus elephantum (Hance) A.Camus		x		3/6	732041	1712500
69	Fagaceae	Lithocarpus pierrei (Hickel & A.Camus) A.Camus		x		1/1	752943	1719617
70	Fagaceae	Lithocarpus pierrei (Hickel & A.Camus) A.Camus		x		1/6	752607	1719693
71	Fagaceae	Lithocarpus pierrei (Hickel & A.Camus) A.Camus		x		2/1	750740	1722688
72	Fagaceae	Lithocarpus pierrei (Hickel & A.Camus) A.Camus		x		2/4	752726	1722359
73	Fagaceae	Lithocarpus pierrei (Hickel & A.Camus) A.Camus		x		2/6	752943	1720599
74	Fagaceae	Lithocarpus pierrei (Hickel & A.Camus) A.Camus		x		3/2	735209	1712032
75	Fagaceae	Lithocarpus pierrei (Hickel & A.Camus) A.Camus		x		3/6	732041	1712500
76	Fagaceae	Lithocarpus pierrei (Hickel & A.Camus) A.Camus		x		5/1	723071	1710374
77	Fagaceae	Lithocarpus pierrei (Hickel & A.Camus) A.Camus		x		5/2	723254	1710986
78	Fagaceae	Lithocarpus pierrei (Hickel & A.Camus) A.Camus		x		5/3	722916	1711735
79	Fagaceae	Lithocarpus pierrei (Hickel & A.Camus) A.Camus		x		5/6	722534	1713764
80	Lauraceae	Machilus sp.			x	2/2	750569	1723483
81	Sapotaceae	Madhuca cochinchinensis (Pierre ex Dubard) H.J.Lam		x		2/5	752836	1721953
82	Lauraceae	Neolitsea sp.			x	1/5	751734	1717636
83	Escalloniaceae	Polyosma dolichocarpa Merr.		x		2/4	752726	1722359
84	Escalloniaceae	Polyosma dolichocarpa Merr.		x		2/6	752943	1720599
85	Rubiaceae	Psychotria cambodiana Pierre ex Pit.		x		4/1	728886	1712825
86	Rubiaceae	Psychotria cambodiana Pierre ex Pit.		x		4/2	728974	1712562
87	Rubiaceae	Psychotria cambodiana Pierre ex Pit.		x		4/3	729218	1712797
88	Fagaceae	Quercus langbianensis Hickel & A.Camus	NT	x		4/1	728886	1712825
89	Fagaceae	Quercus langbianensis Hickel & A.Camus	NT	x		4/2	728974	1712562
90	Fagaceae	Quercus langbianensis Hickel & A.Camus	NT	x		4/3	729218	1712797
91	Fagaceae	Quercus langbianensis Hickel & A.Camus	NT	x		4/4	730017	1712249
92	Fagaceae	Quercus langbianensis Hickel & A.Camus	NT	x		4/5	730445	1712019
93	Fagaceae	Quercus sp.1			x	1/1	752943	1719617
94	Fagaceae	Quercus sp.1			x	1/2	752607	1718678
95	Sapotaceae	Sarcosperma kontumense Gagnep. ex Aubrév.		x		2/4	752726	1722359
96	Symplocaceae	Symplocos wikstroemiifolia Hayata		x		4/1	728886	1712825
97	Symplocaceae	Symplocos wikstroemiifolia Hayata		x		4/2	728974	1712562
98	Symplocaceae	Symplocos wikstroemiifolia Hayata		x		4/3	729218	1712797
99	Symplocaceae	Symplocos wikstroemiifolia Hayata		x		4/4	730017	1712249
100	Symplocaceae	Symplocos wikstroemiifolia Hayata		x		4/5	730445	1712019
101	Symplocaceae	Symplocos wikstroemiifolia Hayata		x		4/6	731205	1712548
102	Symplocaceae	Symplocos wikstroemiifolia Hayata		x		5/1	723071	1710374
103	Symplocaceae	Symplocos wikstroemiifolia Hayata		x		5/6	722534	1713764
104	Rubiaceae	Urophyllum sp. 1 (new genus record for Laos)		x		2/1	750740	1722688

106	Rubiaceae	<i>Urophyllum</i> sp. 3 (new genus record for Laos)		x		2/6	752943	1720599
107	Ericaceae	<i>Vaccinium</i> sp. 1			x	2/2	750569	1723483
108	Polygalaceae	<i>Xanthophyllum ellipticum</i> Korth. ex Miq.		x		1/3	752318	1717523
109	Polygalaceae	<i>Xanthophyllum ellipticum</i> Korth. ex Miq.		x		1/5	751734	1717636
110	Polygalaceae	<i>Xanthophyllum ellipticum</i> Korth. ex Miq.		x		2/1	750740	1722688
111	Polygalaceae	<i>Xanthophyllum ellipticum</i> Korth. ex Miq.		x		2/2	750569	1723483
112	Polygalaceae	<i>Xanthophyllum ellipticum</i> Korth. ex Miq.		x		2/3	751025	1722672
113	Polygalaceae	<i>Xanthophyllum ellipticum</i> Korth. ex Miq.		x		2/4	752726	1722359
114	Polygalaceae	<i>Xanthophyllum</i> sp. 1			x	5/1	723071	1710374
115	Zingiberaceae	<i>Zingiber mellis</i> Škorničk., H.B.Trần & Šída f.	EN	x		3/3	735792	1712515
116	Podocarpaceae	<i>Nageia fleuryi</i> (Hickel) de Laub.	NT			2/3	751025	1722672
117	Podocarpaceae	<i>Nageia fleuryi</i> (Hickel) de Laub.	NT			5/6	722534	1713764
118	Pinaceae	<i>Pinus dalatensis</i> Ferré	NT			2/1	750740	1722688
119	Pittosporaceae	<i>Pittosporum pauciflorum</i> Hook. & Arn.	VU			4/1	728886	1712825
120	Pittosporaceae	<i>Pittosporum pauciflorum</i> Hook. & Arn.	VU			4/6	731205	1712548
121	Pittosporaceae	<i>Pittosporum pauciflorum</i> Hook. & Arn.	VU			5/1	723071	1710374
122	Pittosporaceae	<i>Pittosporum pauciflorum</i> Hook. & Arn.	VU			5/2	723254	1710986
123	Pittosporaceae	<i>Pittosporum pauciflorum</i> Hook. & Arn.	VU			5/3	722916	1711735

**Mammals: Coordinates of Globally Threatened Species**

Group and Family	Common Name	Scientific Name	GT	Endemic Species			Survey block	UTM		Alt (m a.s.l)
				First Record	Second Record	New SP Can.		X	Y	
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				2	751472	1722999	1,259
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				3	722825	1712721	1,340
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				2	753085	1722328	1,140
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				3	732291	1711333	1,598
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				2	752030	1720813	1,209
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				2	753233	1721151	1,095
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				2	752624	1722764	1,220
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				3	733417	1712501	1,494
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				3	732425	1712541	1,599
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				3	733458	1711473	1,517
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				4	730566	1709941	1,559
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				4	729860	1709973	1,467
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				4	729097	1711619	1,311
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				4	731020	1711408	1,615
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				4	730566	1709941	1,559
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				4	729097	1711619	1,311
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				5	721848	1711560	1,219
<i>Colobinae</i>	Stump-tailed Macaque	<i>Macaca arctoides</i>	VU				5	721100	1711389	1,206
<i>Hylobatinae</i>	Buff-cheeked Gibbon	<i>Nomascus annamensis</i>	EN				1	753461	1719462	1,135
<i>Hylobatinae</i>	Buff-cheeked Gibbon	<i>Nomascus annamensis</i>	EN				1	754059	1719265	1,167
<i>Hylobatinae</i>	Buff-cheeked Gibbon	<i>Nomascus annamensis</i>	EN				2	753416	1721669	1,160
<i>Hylobatinae</i>	Buff-cheeked Gibbon	<i>Nomascus annamensis</i>	EN				4	729900	1710465	1,439
<i>Hylobatinae</i>	Buff-cheeked Gibbon	<i>Nomascus annamensis</i>	EN				4	729956	1711647	1,510
<i>Colobinae</i>	Red-shanked Douc Langur	<i>Pygathrix nemaus</i>	CR				2	751402	1722656	1,243
<i>Colobinae</i>	Red-shanked Douc Langur	<i>Pygathrix nemaus</i>	CR				3	732493	1712583	1,623
<i>Colobinae</i>	Red-shanked Douc Langur	<i>Pygathrix nemaus</i>	CR				4	730836	1710015	1,600
<i>Colobinae</i>	Red-shanked Douc Langur	<i>Pygathrix nemaus</i>	CR				4	730837	1710699	1,565
<i>Colobinae</i>	Red-shanked Douc Langur	<i>Pygathrix nemaus</i>	CR				4	730837	1710015	1,600
<i>Colobinae</i>	Red-shanked Douc Langur	<i>Pygathrix nemaus</i>	CR				4	730838	1710700	1,565
<i>Colobinae</i>	Red-shanked Douc Langur	<i>Pygathrix nemaus</i>	CR				4	730085	1710377	1,339
<i>Colobinae</i>	Red-shanked Douc Langur	<i>Pygathrix nemaus</i>	CR				4	730566	1709941	1,559
<i>Manidae</i>	Chinese Pangolin	<i>Manis pentadactyla</i>	CR				1	751113	1717914	1,133
<i>Manidae</i>	Chinese Pangolin	<i>Manis pentadactyla</i>	CR				1	751078	1717929	1,123
<i>Manidae</i>	Chinese Pangolin	<i>Manis pentadactyla</i>	CR				1	751090	1717921	1,129
<i>Manidae</i>	Chinese Pangolin	<i>Manis pentadactyla</i>	CR				2	750637	1724419	1,210
<i>Manidae</i>	Chinese Pangolin	<i>Manis pentadactyla</i>	CR				2	751037	1723050	1,107
<i>Manidae</i>	Chinese Pangolin	<i>Manis pentadactyla</i>	CR				4	729628	1709538	1,357
<i>Manidae</i>	Chinese Pangolin	<i>Manis pentadactyla</i>	CR				4	728728	1709978	1,157
<i>Manidae</i>	Sunda Pangolin	<i>Manis javanica</i>	CR				2	752546	1722746	1,117
<i>Manidae</i>	Chinese Pangolin	<i>Manis pentadactyla</i>	CR				2	753134	1721101	1,081
<i>Manidae</i>	Chinese Pangolin	<i>Manis pentadactyla</i>	CR				2	751361	1722498	1,249
<i>Manidae</i>	Sunda Pangolin	<i>Manis javanica</i>	CR				2	750623	1723932	1,240
<i>Manidae</i>	Sunda Pangolin	<i>Manis javanica</i>	CR				2	752546	1722746	1,117
<i>Manidae</i>	Sunda Pangolin	<i>Manis javanica</i>	CR				1	753904	1718763	1,103
<i>Manidae</i>	Sunda Pangolin	<i>Manis javanica</i>	CR				1	751740	1717708	1,077
<i>Manidae</i>	Sunda Pangolin	<i>Manis javanica</i>	CR				1	751875	1718740	1,096
<i>Manidae</i>	Sunda Pangolin	<i>Manis javanica</i>	CR				1	750995	1717133	1,055
<i>Manidae</i>	Sunda Pangolin	<i>Manis javanica</i>	CR				1	751718	1717678	1,055
<i>Manidae</i>	Sunda Pangolin	<i>Manis javanica</i>	CR				1	751740	1717708	1,055
<i>Manidae</i>	Sunda Pangolin	<i>Manis javanica</i>	CR				1	751652	1718019	1,108
<i>Manidae</i>	Sunda Pangolin	<i>Manis javanica</i>	CR				2	752645	1720851	1,135
<i>Ursidae</i>	Asiatic Black Bear	<i>Ursus thibetanus</i>	VU				2	751402	1722656	1,220
<i>Ursidae</i>	Sun Bear	<i>Helarctos malayanus</i>	VU				2	752479	1722669	1,226
<i>Ursidae</i>	Sun Bear	<i>Helarctos malayanus</i>	VU				2	751325	1722243	1,248
<i>Ursidae</i>	Sun Bear	<i>Helarctos malayanus</i>	VU				4	729339	1710074	1,255
<i>Ursidae</i>	Sun Bear	<i>Helarctos malayanus</i>	VU				4	728770	1709947	1,184
<i>Ursidae</i>	Sun Bear	<i>Helarctos malayanus</i>	VU				4	728834	1709911	1,212

Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			4	728838	1709910	1,230
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723129	1712261	1,284
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723266	1712237	1,287
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723382	1712245	1,279
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723520	1712263	1,273
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723520	1712262	1,268
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723564	1712247	1,243
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723564	1712254	1,251
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723643	1712291	1,267
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723642	1712298	1,270
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723665	1712334	1,231
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	721642	1713606	1,323
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723266	1712237	1,287
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723382	1712245	1,279
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723520	1712263	1,273
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723520	1712262	1,268
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723564	1712247	1,243
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723564	1712254	1,251
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723643	1712291	1,267
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723642	1712298	1,270
Ursidae	Sun Bear	<i>Helarctos malayanus</i>	VU			5	723665	1712334	1,231
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	732425	1712541	1,599
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	731020	1711408	1,615
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	732923	1712124	1,574
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	729097	1710186	1,205
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	732923	1712124	1,574
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	729934	1710026	1,450
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730190	1710041	1,455
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730197	1710040	1,455
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730303	1709908	1,510
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730303	1709897	1,510
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730672	1711061	1,595
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730967	1711228	1,617
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730969	1711206	1,633
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730975	1711190	1,630
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730977	1711342	1,355
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730977	1711343	1,590
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			5	723011	1713638	1,307
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			2	751570	1720922	1,218
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			2	751361	1720881	1,228
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			2	749913	1722013	1,175
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			2	749758	1721516	1,202
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	734177	1711333	1,314
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	733865	1710656	1,446
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	733556	1710165	1,333
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	733292	1709346	1,377
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	732551	1710388	1,344
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	732681	1711965	1,667
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	732758	1711932	1,671
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	732815	1711948	1,677
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	732819	1711947	1,677
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	732982	1712151	1,536
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	733362	1709377	1,386
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	733910	1709537	1,372
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			3	733342	1709203	1,359
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	729021	1711816	1,220
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	729022	1710187	1,191
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	729002	1710199	1,193
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	728857	1710251	1,223
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	728435	1710597	1,212
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	727902	1710831	1,252
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730788	1710614	1,559
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	730900	1710863	1,604
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	729762	1710465	1,277
Bovidae	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU			4	728837	1709926	1,219

<i>Bovidae</i>	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU				4	728838	1709926	1,219
<i>Bovidae</i>	Chinese Serow	<i>Capricornis milneedwardsii</i>	VU				4	728966	1710048	1,201
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				5	721100	1711389	1,206
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				1	751361	1722393	1,243
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				1	751357	1722333	1,247
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				1	751999	1716192	1,015
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				2	751357	1722333	1,247
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				3	734416	1711606	1,244
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				3	734540	1711897	1,234
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				1	752165	1716093	1,126
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				1	753475	1718680	1,105
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				1	751873	1718713	1,081
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				2	749864	1722653	1,155
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				2	749874	1722628	1,154
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				2	749861	1722258	1,157
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				2	750719	1722436	1,255
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				2	750835	1722538	1,262
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				2	750719	1722436	1,233
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				2	750835	1722538	1,242
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				3	734364	1713062	1,242
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				4	729375	1710315	1,276
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				4	729543	1710458	1,276
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				4	729726	1710575	1,300
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				4	729799	1711609	1,314
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				4	729630	1710450	1,272
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				5	721483	1713559	1,285
<i>Cervidae</i>	Sambar Deer	<i>Cervus unicolor</i>	VU				5	721503	1713582	1,297
<i>Mustelidae</i>	Smooth-coated Otter	<i>Lutrogale perspicillata</i>	VU				3	732896	1709422	1,341
<i>Mustelidae</i>	Great Hog Badger	<i>Arctonyx collaris</i>	VU				5	721472	1713389	1,324
<i>Cercopithecidae</i>	Northern Pig-tailed Macaque	<i>Macaca leonina</i>	VU				5	721848	1711560	1,219
<i>Cercopithecidae</i>	Northern Pig-tailed Macaque	<i>Macaca leonina</i>	VU				5	721848	1711560	1,219
<i>Viverridae</i>	Owston's Civet	<i>Chrotogale owstoni</i>	EN				3	733458	1711473	1,517

1 Herps: Coordinates of first and second records, Globally Threatened and New Species Candidates										
2	Group and Family	Species	GT	Endemism Species			Survey block	Coordinates		Alt (m a.s.l)
				First Record	Second Record	New SP Candidate		N	E	
4	<b>LIZARDS</b>									
5	Anguidae	Dopasia sokolovi			x		4	15.470230	107.182620	1322
6	Agamidae	Acanthosaura prasina		x			4	15.469810	107.175920	1520
7	Scincidae	Pseudocalotes ziegleri			x		3	15.478276	107.154488	1440
8	Scincidae	Plestiodon quadrilineatus		x			1	15.535272	107.338804	1080
9	Elaphidae	Bungarus slowinskii	VU				2	15.555660	107.358260	1080
10										
11	<b>AMPHIBIANS – FROGS</b>									
12	Ranidae	Amolops spinaepectoralis		x			3	15.502398	107.368455	1170
13	Dicroglossidae	Quasipaa sp.				x		15.478276	107.154488	1440
14	Dicroglossidae	Limnonectes poilani		x				15.539671	107.353805	1065
15	Megophryidae	Leptobrachella firthi		x			1	15.544283	107.352045	1080
16	Megophryidae	Xenophrys cf maosonensis				x	2	15.544283	107.352045	1090
17										
18	Rhacophoridae	Rhacophorus sp nov				x	2	15.459712	107.285291	1185
19	Rhacophoridae	Theioderma truongsoneensis		x			4	15.478276	107.154488	1455
20	Rhacophoridae	Kurixalus cf gryllus	VU				4	15.478428	107.152935	1450

## Annex 7. Some pictures of wildlife from camera trapping



Cam 1\_SB1: Annamite Muntjac



Cam 1\_SB1: Small-toothed Ferret Badger



Cam 1\_SB1: Silvered Pheasant (female)



Cam 1\_SB1: Masked Palm Civet



Cam 2\_SB1: East Asian Porcupine



Cam 2\_SB1: Annamite Muntjac



Cam 3\_SB1: Silvered Pheasant (female)



Cam 3\_SB1: Berdmores berylmys Rat



Cam 5\_SB1: Wild Pig



Cam 1\_SB2: Wild Pig (?)



Cam 2\_SB2: Red Muntjac



Cam 2\_SB2: Masked Palm Civet



Cam 2\_SB2: Wild Pig (?)



Cam 2\_SB2: Annamite Muntjac



Cam 2\_SB2: Red Muntjac




Cam 2\_SB2: Annamite Muntjac



Cam 2\_SB2: Large Indian Civet



Cam 2\_SB2: Eurasian Wild Pig



Cam 3\_SB2: Stump-tailed Macaque (VU)



Cam 3\_SB2: Annamite Muntjac



Cam 3\_SB2: Black-hood Laughingthrush



Cam 3\_SB2: Red Muntjac



Cam 3\_SB2: Wild Pig (?)



Cam 3\_SB2: Silvered Pheasant (sub-species) – male



Cam 4\_SB2: Annamite Muntjac



Cam 4\_SB2: Stump-tailed Macaque (VU)



Cam 4\_SB2: Wild Pig (?)



Cam 5\_SB2: Silvered Pheasant (sub-species) – male



Cam 6\_SB2: Wild Pig



Cam 6\_SB2: Stump-tailed Macaque (VU)



Cam 2\_SB3: Stump-tailed Macaque (VU)



Cam 2\_SB3: Annamite Muntjac



Cam 2\_SB3: Chinese Serow (VU)



Cam 3\_SB3: Crab-eating Mongoose



Cam 3\_SB3: Stump-tailed Macaque (VU)



Cam 4\_SB3: Chinese Serow (VU)



Cam 5\_SB3: Brush-tailed Porcupine



Cam 5\_SB3: Silvered Pheasant (male)



Cam 1\_SB4: Stump-tailed Macaque (VU) Cam 1\_SB4: Crab-eating Mongoose



Cam 1\_SB4: Chinese Serow (VU)



Cam 2\_SB4: Annamite Muntjac



Cam 2\_SB4: Stump-tailed Macaque (VU)



Cam 2\_SB4: Red Muntjac



Cam 3\_SB4: Annamite Muntjac



Cam 3\_SB4: Silvered Pheasant (males and female)



Cam 3\_SB4: Red-shanked Douc Langur (CR)



Cam 3\_SB4: Red Muntjac



Cam 3\_SB4: Stump-tailed Macaque (VU)



Cam 3\_SB4: Small Asian Mongoose



Cam 3\_SB5: Northern Pig-tailed Macaque (VU)



Cam 4\_SB4: Brush-tailed Porcupine



Cam 4\_SB4: Annamite Muntjac (female?)



Cam 4\_SB4: Stump-tailed Macaque (VU)



Cam 4\_SB4: Annamite Muntjac



Cam 4\_SB4: Crab-eating Mongoose




Cam 4\_SB4: Masked Palm Civet



Cam 5\_SB4: Long-tailed Giant Rat



Cam 5\_SB4: Chinese Serow (VU)



Cam 5\_SB4: Small-toothed Ferret Badger Marten

Cam 5\_SB4: Yellow-throated



Cam 5\_SB4: Annamite Muntjac

Cam 5\_SB4: Eurasian Wild Pig



Cam 6\_SB4: Silvered Pheasant (males, sub-species)

Cam 6\_SB4: Annamite Muntjac



Cam 1\_SB5: Yellow-throated Marten



Cam 1\_SB5: Common Treeshrew (?)



Cam 1\_SB5: Annamite Muntjac



Cam 1\_SB5: Small-toothed Ferret Badger



Cam 1\_SB5: Spotted Linsang

Cam 1\_SB5: Long-tailed Giant Rat



Cam 1\_SB5: Masked Palm Civet

Cam 1\_SB4: Pallas's squirrel



Cam 1\_SB5: Variable Squirrel (?)



Cam 2\_SB5: Yellow-throated Marten



Cam 2\_SB5: Annamite Muntjac (?)



Cam 2\_SB5: Northern Pig-tailed Macaque (VU)



Cam 2\_SB5: Red Muntjac



Cam 2\_SB5: Sambar Deer (VU)



Cam 3\_SB5: Black Giant Squirrel



Cam 3\_SB5: Annamite Muntjac



Cam 3\_SB5: Red Muntjac



Cam 3\_SB5: Red Junglefowl



Cam 4\_SB5: Annamite Muntjac



Cam 4\_SB4: Stump-tailed Macaque (VU)



Cam 5\_SB5: Wild Pig



Cam 6\_SB5: Great Hog Badger (VU)



Cam 5\_SB5: Annamite Muntjac



Cam 5\_SB3: Annamite Muntjac



Cam 5\_SB3: Crab-eating Mongoose



Cam 6\_SB3: Owston's Civet  
Badger



Cam 6\_SB3: Masked Palm Civet



Cam 6\_SB3: Small-toothed Ferret



Cam 7\_SB3: Annamite Muntjac

## Annex 8. Some other incidental records

### *Fish*

*Fish*: a total of 9 fish species were found by opportunistic in small upstream of SB1 and SB2 with some photographs, but did not listed for this report. Most interesting was the record of important endemic genus which is usually found only in upstream at high elevation such such as the Genus of *Schistura*, *Annamia*, *Vamanenia* and *Poropuntius* (see below).



A. *Annamia* sp., B. *Vamanenia* sp., C. *Schistura* sp.  
D. *Channa limbata*

### *Bird*

The bird survey was conducted by the bird team separately from this assignment; however, a total of over 130 bird species were recorded by opportunistic encounters with some photographs during the surveys especially from the dry season as more winter birds visiting the area. Most interesting was the record of Crested Argus *Rheinardia ocellata* (EN) which was detected from its morning song was heard in the block 1 on July 13 and 14, 2021. This bird was present along the border area with Vietnam. During the dry season this bird gave a call every day in SB1 which were heard from the Base-Camp 1 on the east and north. Often, its calls were heard in evening at at 6pm and morning at 7am from December 6 to 13, 2021. The Annamite Mountain Range of Lao and Vietnam is the habitat of this bird, but it is really rare. As large ground bird is vulnerable to be trapped and was extirpated in many places of the Annamite. It would be present in SB2 as well but we did not observe in early morning

and evening since we have no sub-camp in this block. Feathure of Crested Argus was found in Ban Dak Ta-ok, it was hunted with snare in 2021 from the SB1 - Lao-Vietnam border (see below).



Feather of Crested Argus from Ban Dak Ta-ok noy

Also, feathers of Silvered Pheasant *Lophura nycthemera* and Siamese Fireback *Lophura diardi* and were recorded in SB1 and SB2. Interestingly, several males and females of Silvered Pheasant were caught on camera traps (SB1, SB2, SB3, and SB4). According to the bird

photos from the camera traps is a sub-species to Silvered Pheasant. It is rare and first record of Laos.

As well as Wreathed Hornbill *Aceros nipalensis* (VU) as 5 individuals were seen directly in SB1 from flying overhead during the dry season survey on December 8, 2021, just near the Lao-Vietnam border; and Great Hornbill *Buceros bicornis* (VU) as a pair was seen flying overhead in SB2 on December 10, 2021. The Wreathed Hornbill is the first record in the area and not listed in the IBAT for the area. These birds are nationally conservation significance. In addition, other important and endemic birds are Oriental Pied Hornbill *Anthracoceros albirostris*, White-stripped Magpie, Vietnamese Cutia *Cutia legalleni*.

Seasonal variation and distribution of bird species in the survey area is quite different as more species of birds were observed during the dry season since more winter birds visiting the area, and some specific birds were observed such as Wreathed Hornbill (VU) and White-stripped Magpie which were not observed during the wet season survey.



ສາທາລະນະລັດ ປະຊາທິປະໄຕ ປະຊາຊົນລາວ  
ສັນຕິພາບ ເອກະລາດ ປະຊາທິປະໄຕ ເອກະພາບ ວັດທະນະຖາວອນ

ແຂວງເຊກອງ  
ພະແນກກະສິກໍາ ແລະ ປ່າໄມ້

2047  
ເລກທີ: \_\_\_/ກປຂ.21  
ເຊກອງ, ວັນທີ: 30.11.2021

**ໃບສະເໜີ**

ຮຽນ: ຫົວໜ້າຄະນະສະເພາະກິດ ຄວບຄຸມພະຍາດ COVID-19 ຂັ້ນແຂວງ ທີ່ນັບຖື  
ເລື່ອງ: ຂໍສະເໜີທ່ານ ອອກຂໍ້ຕົກລົງແຕ່ງຕັ້ງພະນັກງານລົງເຂັດວຽກປູຜາກສະໜາມທີ່ເມືອງດາກຈິງ  
ແຂວງ ເຊກອງ

- ອີງຕາມ: ຄໍາສັ່ງເນະນໍາ ສະບັບເລກທີ 42/ຈຂ.ຊກ, ລົງວັນທີ 20.8. 2021 ວ່າດ້ວຍສືບຕໍ່ການປ້ອງກັນ ແລະ ຄວບຄຸມຕ້ານ ແລະ ສະກັດກັ້ນການລະບາດຂອງພະຍາດ ໂຄວິດ-19 ທົ່ວແຂວງເຊກອງ.
- ອີງຕາມ: ຫົງສີສະເໜີ ຂອງທີມງານຊ່ຽວຊານ ຈາກມະຫາວິທະຍາໄລແຫ່ງຊາດ ໃນຄັ້ງວັນທີ 23 /11/ 2021 ເລື່ອງການລົງສໍາຫຼວດຊີວະນາໆພັນຂອງທີມງານ ໃນນາມຊ່ຽວຊານຂອງບໍລິສັດ Impact Energy Asia Developmnt Limited (IEAD) ໃນໂຄງການພະລັງງານລົມ ທີ່ເມືອງດາກຈິງ, ແຂວງ ເຊກອງ ລະຫວ່າງວັນທີ 28 ພະຈິກ ຫາວັນ ທີ 20 ທັນວາ 2021

ພະແນກກະສິກໍາ ແລະ ປ່າໄມ້ແຂວງ ຂໍຖືເປັນກຽດຮຽນສະເໜີມາຍັງ ຄະນະສະເພາະກິດຂັ້ນແຂວງ ເພື່ອຕົ້ນຄວ້າ ແລະ ພິຈາລະນາອະນຸມັດ ໃຫ້ຄະນະທີມງານຊ່ຽວຊານ ໃນການລົງສໍາຫຼວດຊີວະນາໆພັນ ໃນນາມຊ່ຽວຊານຂອງບໍລິສັດ Impact Energy Asia Developmnt Limited (IEAD) ໃນໂຄງການພະລັງງານລົມ ທີ່ເມືອງດາກຈິງ, ແຂວງເຊກອງ ລະຫວ່າງ ວັນທີ 28 ພະຈິກ ຫາ ວັນທີ 20 ທັນວາ 2021 ເຊິ່ງມີລາຍຊື່ດັ່ງລຸ່ມນີ້:

- ພາກສ່ວນທີມງານຊ່ຽວຊານຈາກມະຫາວິທະຍາໄລແຫ່ງຊາດ ມີຈຳນວນ 11 ຄົນ :
  1. ທ່ານ ປອ ໄພວັນ ເພຍປະລັດ
  2. ທ່ານ ປອ ທະນິນ ໂຄດປະທຸມ
  3. ທ່ານ ປອ ເພັດລາສີ ສຸລະເດັດ
  4. ທ່ານ ປອ ມັດມະນີ ສຸຂະວິງ
  5. ທ່ານ ດວງພະຈິນ ສຸວັນໄຊ
  6. ທ່ານ ທອງ ວົງພູທອນ
  7. ທ່ານ ບິດເຕີ ເຊເກວ
  8. ທ່ານ ນາທາແນວ ມິວລີ
  9. ທ່ານ ສຸນທອນ ພິລາວິງ
  10. ທ່ານ ແນຍ ທອນພິກດີ
  11. ທ່ານ ວິລາສັກ ຈັນທະບົວສອນ
- ພາກສ່ວນພະນັກງານວິຊາການຂັ້ນແຂວງ 1 ທ່ານ :
  1. ທ່ານ ສິມຈິດ ບຸລະພອນ ຫົວໜ້າໜ່ວຍງານລົງສັດ, ຂະແໜງລົງສັດ ແລະ ການປະມົງ ດັ່ງນັ້ນ ຈຶ່ງຮຽນສະເໜີມາຍັງທ່ານເພື່ອຕົ້ນຄວ້າ ແລະ ພິຈາລະນາຕາມຄວາມສົມຄວນນີ້ດ້ວຍ. ๗

ຫົວໜ້າພະແນກກະສິກໍາ ແລະ ປ່າໄມ້ ✓  


ເສີມສີ ສຸລິຕະ

## Annex 10. List of External Experts for Consultations

<b>Name of Expert</b>	<b>Field of expertise</b>	<b>Organization</b>
Dr. Mark Newman	Botany	Royal Botanic Garden Edinburgh, Scotland
Dr. Philip Thomas	Botany	Royal Botanic Garden Edinburgh, Scotland
Dr. Shuichiro Tangane	Botany	Kagoshima University, Japan
Dr. Tetsukazu Yahara	Botany	Kyushu Open University, Japan
Dr. Ngoc Nguyen Van	Botany	Da Lat University, Vietnam
Dr. Somran Suddee	Botany	Forest Herbarium, Thailand
Dr. Chatchai Ngernsaengsaruy	Botany	Kasetsart University, Thailand
Dr. Stuart Lindsay	Botany	Singapore Botanic Garden, Singapore
Dr. Will Duckworth	Wildlife	SCC of IUCN HQ/ Senior Biologist of WCS
Dr. Rob Timmins	Wildlife	Senior Biologist of WCS/Independent Researcher
Dr. Duc Hoang Minh	Wildlife	Southern Institute of Ecology, Ho Chi Minh
Dr. Bryan Stuart	Reptile	North Carolina Museum of Natural Sciences
Dr. Somphouthone Phimmachak	Reptile	National University of Laos, Vientiane



# **COLLISION RISK MODEL FOR ESTIMATING ANNUAL MORTALITY OF GREY-FACED BUZZARD (BUTASTUR INDICUS)**

[Double-click to import picture]



## Collision Risk Model for IEAD Monsoon Wind Power Project

Estimating annual mortality of Grey-faced  
buzzard (*Butastur indicus*)

15th February 2022

Project No.: 0598121

Document details	
Document title	Collision Risk Model for IEAD Monsoon Wind Power Project
Document subtitle	Estimating annual mortality of Grey-faced buzzard ( <i>Butastur indicus</i> )
Project No.	0598121
Date	15th February 2022
Version	Draft 0.02
Author	Sebastian Ellis
Client Name	IEAD

#### Document history

Version	Revision	Author	Reviewed by	ERM approval to issue		Comments
				Name	Date	
Draft	01	Sebastian Ellis	Les Hatton	Name	00.00.0000	First draft
Draft	02	Sebastian Ellis				Comments addressed, location figure added

---

**Signature Page**

15th February 2022

# Collision Risk Model for IEAD Monsoon Wind Power Project

Estimating annual mortality of Grey-faced buzzard (*Butastur indicus*)

[Double click to insert signature]

Name  
Job title

[Double click to insert signature]

Name  
Job title

[Double click to insert signature]

Name  
Job title

[Double click to insert signature]

Name  
Job title

Environmental Resources Management Limited  
2<sup>nd</sup> Floor, Exchequer Court  
33 St Mary Axe  
London  
EC3A 8AA

© Copyright 2022 by ERM Worldwide Group Ltd and/or its affiliates ("ERM").  
All rights reserved. No part of this work may be reproduced or transmitted in any form,  
or by any means, without the prior written permission of ERM.

## CONTENTS

<b>1. INTRODUCTION .....</b>	<b>5</b>
1.1 Purpose and scope.....	5
<b>2. METHODOLOGY AND DATA .....</b>	<b>5</b>
2.1 Excel spreadsheet calculations.....	5
2.2 Collision Risk Model .....	5
2.2.1 Bird Data.....	5
2.2.2 Turbine and Array Data.....	7
2.2.3 Collision Calculations.....	9
2.3 Potential Biological Removal .....	11
<b>3. RESULTS .....</b>	<b>12</b>
3.1 Estimated Annual Mortality of Grey-faced buzzard.....	12
3.2 Potential Biological Removal of Grey-faced buzzard .....	13
<b>4. SUMMARY .....</b>	<b>14</b>
4.1 Interpreting Estimated Annual Mortality and Potential Biological Removal of Grey-faced buzzard.....	14
<b>5. REFERENCES .....</b>	<b>15</b>

### List of Tables

Table 1: Bird data used within CRM .....	6
Table 2: VP Survey Data for Each Array .....	6
Table 3: Turbine array data used within CRM .....	8
Table 4: Turbine technical specifications used within CRM.....	8
Table 5: Risk of Collision of a Grey-faced buzzard with a turbine .....	9
Table 6: Example calculation of bird occupancy at Ban Dakdongna array .....	10
Table 7: Estimated Annual Mortality – 95% Avoidance .....	13
Table 8: Estimated Annual Mortality – 0% Avoidance .....	13
Table 9: Collision Risk Percentage turbine comparison .....	13
Table 10: Potential Biological Removal for different estimated Grey-face buzzard populations .....	14
Table 11: Impacts of Estimated Annual Mortality and Potential Biological Removal.....	14

### List of Figures

Figure 1: Locations of arrays and VPs within the arrays ..... **Error! Bookmark not defined.**

### Acronyms and Abbreviations

CRH	Collision risk height
CRM	Collision risk model
NatureScot	Scotland's nature agency
PBR	Potential biological removal
RPM	Revolutions per minute
VP	Vantage point
WTG	Wind turbine generator

## 1. INTRODUCTION

### 1.1 Purpose and scope

This report follows from the Monsoon Wind Power Project, Sekong and Attapue Provinces, Lao PDR - Collision Risk Approach report and contains the methods and results of Collision Risk Modelling (CRM) and Potential Biological Removal (PBR) calculations, specifically in regards to the Grey-faced buzzard (*Butastur indicus*).

Grey-faced buzzard was chosen as a species of interest for this modelling based on analysis of datasheets and reports from Dr Santi Xayyasith and his bird survey team. During vantage point (VP) surveys it was encountered at 11 VPs of a total 14 and assessed to have been flying at collision risk height (CRH) for a total of 2190 seconds at 8 VPs within areas of proposed wind turbine generator (WTG) arrays.

To assess the potential impact of the Monsoon Wind Power Project on the Grey-faced buzzard population, collision risk modelling to estimate annual mortality as a result of the project was undertaken following the Band onshore model outlined in the Wind farm impacts on birds - Calculating a theoretical collision risk assuming no avoiding action guidance note (NatureScot, 2000) and a potential biological removal calculation originally developed for marine mammals (Wade, 1998) and since adopted for estimating sustainable levels of bird mortality (Dillingham and Fletcher, 2008) was undertaken.

Estimated annual mortality was calculated at each of the 6 individual turbine arrays and aggregated to give a total overall estimated annual mortality for the project.

## 2. METHODOLOGY AND DATA

### 2.1 Excel spreadsheet calculations

All calculations were made on an excel spreadsheet and rounded to either 1, 2 or 3 decimal places in this report. As such text in figures may not exactly equate to the detailed spreadsheet numbers, due to rounding.

### 2.2 Collision Risk Model

To estimate annual mortality, the collision risk model uses the Band two stage calculation. The calculation first assesses the probability of a bird being hit whilst flying through the rotors, and then secondly applies this probability to the annual number of birds transiting the rotors within a windfarm array. Data used in the model includes bird morphological measurements and physical turbine parameters. Where this data is unknown or in a range, in order to provide a conservative estimate, the 'worst case' numbers are chosen or a reasonable assumption based on similar data is made.

#### 2.2.1 Bird Data

Grey-faced buzzard are a migratory species and they are assumed to have presence for 5 months of the year between November to March, based on Vantage Point data, a total of 151 days. Available daylight hours during the months the grey-faced buzzard is present within the project site averaged 11.55 per day (Timeanddate.com, 2022). As a diurnal species roosting at night this value was used in the model as the hours per day birds were present and flying. Physically grey-faced buzzards are 41-48cm long with a wingspan of 101-110cm (Ferguson-Lees & Christie, 2001), to obtain 'worst case' results 48cm and 110cm were the values used within the model. Very little is known about their flight speed and there is no data available. The 12.5m/s flight speed data of the Harris Hawk (*Parabuteo*

*unicinctus*) is used within the model as a proxy (AZ Animals, 2022), this choice is due to similarity of physical size and behaviour of both species. Within the model there is an option to select flapping or gliding flight type, flapping was chosen as a 'worst case' and due to observed behaviour of raptors interacting/avoiding wind turbines. Grey-faced buzzard avoidance rates with wind turbines are unknown, NatureScot guidance shows raptor avoidance between 95-99% depending on species, as such 95% avoidance rate was chosen as a worst case and applied to the overall estimated annual mortality. These parameters as used in the model are shown in table 1.

Data from Dr Santi Xayyasith's VP datasheets is summarised in table 2 and locations in figure 1. Data from VPs 9 and 10 are not used within the model as these VPs are within an area for a High Voltage cable with no turbines and therefore no turbine collision risk. The single below collision risk height flight recorded from VP 14 in June is likely to be either a species mis-identification or an aberrant record of a wandering none (or failed) breeder. As a consequence of this and the flight being below collision risk height it has been excluded from analysis. The remaining VP data was separated according to the relevant array and was used to calculate risk for each individual array where grey-faced buzzard was recorded.

**Table 1: Bird data used within CRM**

Bird Data	Value
Length (metres)	0.48
Wingspan (metres)	1.1
Flight speed (metres/sec)	12.5*
Days per year bird present (days)	151
Hours per day bird present (hours)	11.55
Avoidance rate (%)	95%

\*Proxy value from Harris Hawk

**Table 2: VP Survey Data for Each Array**

VP	Array	Total Time Surveyed (hours)	Area Visible from VP (hectare)	Hectare hours (time x area)	Total Flight time observed (seconds)	Flight time observed 0-30m (seconds)	Flight time observed 30-150m (seconds)	Flight time observed >150m (seconds)	Data used in CRM calculations (Yes/No)
1	East Central Array	120	628.5	75420	0	0	0	0	Yes
2	East Central Array	120	628.5	75420	484	0	330	135	Yes
3	East Central Array	120	628.5	75420	90	60	30	0	Yes
4	East Central Array	120	628.5	75420	330	135	195	0	Yes
5	South West Array	120	628.5	75420	390	0	345	45	Yes
6	Ban Dakdonna Array	120	628.5	75420	390	195	195	0	Yes

7	South West Array	120	628.5	75420	150	105	45	0	Yes
8	South West Array	120	628.5	75420	0	0	0	0	Yes
9	HV Line	120	628.5	75420	720	45	615	60	No
10	HV Line	120	628.5	75420	300	0	0	300	No
11	Ban Dakdonna Array	120	628.5	75420	0	0	0	0	Yes
12	Dak Cheung village Array	108	628.5	67878	360	105	255	0	Yes
13	North West Array	108	628.5	67878	585	405	180	0	Yes
14	Southernmost Array	108	628.5	67878	30	30	0	0	No



**Figure 1: Locations of the arrays and VPs**

### 2.2.2 Turbine and Array Data

Designs for the project show 155 total turbines split between 6 arrays, the largest array has 44 turbines and the smallest array has 7 turbines. A 500m buffer was created around each turbine in each array in order to calculate total array area as shown in table 3. There are two different turbine designs within the project and parameters used within the model for these turbines were taken from the technical specifications for the GW165-4.0MW and GW155-4.5MW (see table 4). Both turbine designs have 95% availability (aka proportion of time running), 110m hub height above ground level,

and despite being variable pitch by design for the purposes of the CRM they were both assumed to operate at 10 degrees' blade pitch. A collision risk height correction factor was calculated for both turbines to scale up collected survey data from a survey assumed 30m to 150m collision risk height to fit the design collision risk height (27.5m to 192.5m for the GW165-4.0MW and 32.5m to 187.5m for the GW155-4.5MW). The correction factor is calculated by taking the actual size of the turbine collision risk zone (165m for the GW165-4.0MW and 155m for the GW155-4.5MW) and dividing it by the size of the VP survey assumed collision risk zone (120m).

**Table 3: Turbine array data used within CRM**

Array	Number of GW165 – 4.0MW	Number of GW155 – 4.5MW	Number of Total Turbines	Array area (Hectares)
Ban Dakdonna Array	34	3	37	1889.72
Dak Cheung village Array	7	0	7	394.15
East Central Array	34	10	44	2108.36
North West Array	17	0	17	782.00
South West Array	34	7	41	2195.21
Southernmost Array	9	0	9	409.99
Total	135	20	155	7779.44

**Table 4: Turbine technical specifications used within CRM**

Parameter	GW165 – 4.0MW	GW155 – 4.5MW
Total number used in project	135	20
Rotor diameter (metres)	165	155
Rotor blades	3	3
Hub height (metres)	110	110
Minimum tip height above ground level (metres)	27.5	32.5
Maximum tip height above ground level (metres)	192.5	187.5
Maximum blade width (metres)	2.8	4.8
Blade pitch (degrees)	10*	10*
Maximum rated RPM	10.5	9.5
Availability (%)	95%	95%
Collision risk height correction factor**	1.375	1.292

\*Assumed value for CRM, turbine specifications show variable pitch

\*\*VP survey data was collected assuming a collision risk zone between 30m to 150m, this correction factor scales the data up to match actual turbine parameters

## 2.2.3 Collision Calculations

### 2.2.3.1 First Stage – Risk of Collision

The first stage of the CRM calculation uses NatureScot's 'Calculation of collision risk for bird passing through rotor area' spreadsheet (NatureScot, 2000). Inputting turbine and bird parameters yields an upwind, downwind and average collision risk.

Bird length was set as 0.48m and wingspan as 1.1m, with 12.5m/sec flight speed with flapping set as the flight type.

Table 5 shows the parameters used for each turbine design during this first stage of the calculation and the results. Rotation period is defined as time taken for a single rotation and was calculated using turbine maximum rated RPM.

For the GW165-4.0MW turbine average Collision Risk was to 4.02% after 95% wind turbine availability was applied.

For the GW155-4.5MW turbine average Collision Risk was to 5.43% after 95% wind turbine availability was applied.

**Table 5: Risk of Collision of a Grey-faced buzzard with a turbine**

Parameter	GW165 – 4.0MW	GW155 – 4.5MW
Collision risk factoring in turbine availability (%)	4.02	5.43
Collision risk (%)	4.23	5.72
Number of blades	3	3
Maximum width of blade (m)	2.8	4.8
Pitch (degrees)	10	10
Bird length (m)	0.48	0.48
Bird wingspan (m)	1.1	1.1
Flapping (0) or gliding (1)	Flapping (0)	Flapping (0)
Bird speed (m/sec)	12.5	12.5
Rotor diameter (m)	165	155
Rotation period (sec)	5.71	6.32

### 2.2.3.2 Second Stage – Number of transits through rotors

To inform the second stage of the modelling the following calculations were made for each set of turbines at each individual array:

#### *Flight risk volume*

This is defined as the flight risk volume is equal to the maximum height of the rotor (m) multiplied by the area of the array (ha) multiplied by 10,000.

The maximum height of the rotor is taken from the technical specifications as in table 4. The area of the array was the total area of the array divided by the total number of turbines of both types, multiplied by number of the type being calculated as in table 3 (example below). 10,000 is used to convert hectares to metres so the result of the calculation is expressed as metres<sup>3</sup>.

- For example, GW165-4.0MW at Ban Dakdongna array
  - Maximum height: 187.5m
  - Area of the array: 1889.72ha ÷ 37 total turbines × 34 GW165-4.0MW turbines = 1736.5ha
  - Flight risk volume: 3342764749m<sup>3</sup> = 187.5m × (1736.5ha × 10000)

### Combined rotor swept volume

The swept volume is equal to the number of wind turbines multiplied  $\pi R^2$  multiplied by the maximum width of the rotor added to the length of the bird.

The number of the type of wind turbine being calculated in each array is seen in table 3. The radius of each turbine is half the rotor diameter seen in table 4. The maximum width of the blade is seen in table 4 also and the length of the bird is seen in table 1.

- For example, GW165-4.0MW at Ban Dakdongna array
  - Number of GW165-4.0MW turbines in array: 34
  - Radius: 165m ÷ 2 = 82.5m
  - Maximum width of the blade added to length of the bird: 2.8m + 0.48m = 3.28m
  - Combined rotor swept volume: 2384572.497m<sup>3</sup> = 34 ×  $\pi 82.5^2$  × 3.28

### Bird occupancy

Bird occupancy is equal to the number of birds within the array at risk height multiplied by time spent flying in flight risk volume within 1 year.

To calculate the number of birds within the array at risk height from the VP survey data, the total time flight at risk height in hours is divided by the total hectare hours and the result is then multiplied by the total array area in hectares. This gives activity at surveyed risk height across the site. Activity at risk height is then multiplied by collision risk height correction factor for the turbines being calculated (as explained in section 2.2.2 and seen in table 4), giving an adjusted activity at risk height. Adjusted activity at risk height is then multiplied by hours per day that the bird is present on site and days per year the bird is present on site (see table 1) to give bird occupancy.

- For example (see table 6), GW165-4.0MW at Ban Dakdongna array
  - Surveyed activity at risk height: (0.054hours ÷ 150840hectare-hours) × 1889.72hectares = 0.000678599 per hour
  - Adjusted activity at risk height: 0.000678599 × 1.375 = 0.000933074
  - Bird occupancy: 1.627hrs/yr = 0.000933074 × 11.55 × 151

**Table 6: Example calculation of bird occupancy at Ban Dakdongna array**

VP	Array	Total Time Surveyed (hours)	Area Visible from VP (hectare)	Hectare hours (time x area)	Flight time observed 30-150m (hours)
6	Ban Dakdongna Array	120	628.5	75420	0.054
11	Ban Dakdongna Array	120	628.5	75420	0

<b>Total</b>	<b>Total</b>	240	1257	150840	0.054
--------------	--------------	-----	------	--------	-------

### *Bird occupancy of rotor swept volume*

The bird occupancy of rotor swept volume is bird occupancy multiplied by combined rotor swept volume divided by flight risk volume.

Bird occupancy, combined rotor swept volume and flight risk volume have all previously been calculated, there is a multiplication of 3600 to convert the result from hours to seconds.

- For example, GW165-4.0MW at Ban Dakdonna array
  - Bird occupancy of rotor swept volume:  $4.179 = 1.627\text{hrs/yr} \times (2384572.497\text{m}^3 \div 3342764749\text{m}^3) \times 3600$

### *Bird transit time through rotor*

Bird transit time taken is the seconds it takes for a bird to pass through the length of the max rotor width plus bird length.

Using maximum blade width (see table 4), bird length and bird speed (see table 5), it is calculated by adding bird length to blade width and dividing by bird speed.

- For example, GW165-4.0MW at Ban Dakdonna array
  - Bird transit time through rotor:  $0.2624\text{sec} = (2.8\text{m} + 0.48\text{m}) \div 12.5 \text{ m/sec}$

### *Number of transits through rotors*

The number of transits through the rotors is the number of bird expected to fly through the rotors in a year. It is calculated by taking the bird occupancy of the swept rotor volume and dividing it by the bird transit time.

- For example, GW165-4.0MW at Ban Dakdonna array
  - Number of transits through rotors:  $15.926 = 4.179 \div 0.2624$

### *2.2.3.3 Estimated annual number of collisions assuming no avoidance*

Once the above stage one and two calculations are concluded to calculate collision risk with no avoidance the number of transits through rotors in a year is multiplied by the risk of collision factoring in turbine availability (see section 2.2.3.1 and table 5). For the purposes of the model it is assumed all collisions are fatal.

- For example, GW165-4.0MW at Ban Dakdonna array
  - Annual estimated collision risk with no avoidance:  $0.64 = 15.926 \times 4.02\%$

This value is then added to the value obtained for the other turbine design in that array, and summed for all arrays to give an overall estimated annual number of collisions assuming no avoidance.

Avoidance rates as deemed suitable can be applied to this number to get an estimated annual mortality, for this CRM 95% has been chosen as a worst case for raptor avoidance.

## **2.3 Potential Biological Removal**

To calculate PBR the Dillingham and Fletcher (2008) calculation was used.

$$PBR = N_{min} \times \frac{R_{max}}{2} \times F_R$$

$$N_{min} = \tilde{N}e^{(Z_p CV_N)}$$

$$R_{max} = \lambda_{max} - 1$$

$$\lambda_{max} = \frac{(s\alpha - s + \alpha + 1) + \sqrt{(s - s\alpha - \alpha - 1)^2 - 4s\alpha^2}}{2\alpha}$$

This calculation requires input of:

- Estimated population size (individuals) -  $\tilde{N}$
- Recovery factor (0.1-1, where appropriate values may be: 1.0 for 'least concern' species population increasing or stable. 0.5 for 'least concern' species population decreasing. 0.3 for 'near threatened', and 0.1 for all threatened species) -  $F_r$
- Adult survival (0.1-1) -  $s$
- Age at first reproduction (years) -  $\alpha$
- $Z_p$  was set at -0.842 and  $CV_N$  was set at 10% following Dillingham and Fletcher (2008) guidance.

For this assessment of Grey-faced buzzard the following values were used for estimated population. 100,000 individuals as estimated (Ferguson-Lees & Christie, 2001), 50,000 individuals as an assumed new potential number due to population decline, and a worst case scenario of 10,000 individuals at which threshold status on the IUCN Red List may change from Least Concern to Near Threatened. These population numbers were selected as little is known about the true Grey-faced buzzard total population size. The population is generally thought to be decreasing in size due to habitat degradation and loss among other threats (BirdLife International, 2022).

Recovery factor was set as 0.5 for the 100,000 and 50,000 populations, and at 0.3 for the 10,000 as a worst case.

Age at first reproduction was assumed to be 3 years as a worst case, based on a generation length of 5.05 years (BirdLife International, 2022), this also matches with the proxy species of Common Buzzard (*Buteo buteo*).

Adult survival was unknown, for the purposes of the calculation Common Buzzard was chosen as a proxy species with an adult survival of 0.9 (BTO BirdFacts, 2022).

### 3. RESULTS

#### 3.1 Estimated Annual Mortality of Grey-faced buzzard

Following all the calculations detailed in section 2.2.3 based on parameters and data contained in section 2.2 yields the results shown in table 7. To enable other avoidance rates to be applied table 8 contains estimated annual mortality with 0% avoidance.

The overall estimated annual mortality with 95% avoidance of Grey-faced buzzard for the Monsoon Wind Power Project is 0.335 birds per year. Another way to express this is 1 expected Grey-faced buzzard mortality every 3 years. The lowest mortality is expected in the Southernmost array with 0 as during the VP surveys no Grey-faced buzzard were recorded at risk height in this area. The highest mortality is expected in the East Central Array with 0.116, likely due to it having the highest number of turbines of all the arrays.

After factoring in turbine parameters and total percentage on time, the GW155-4.5MW turbine has a slightly higher risk of collision at 5.43% than the GW165-4.0MW with 4.02% (shown in table 9), this is likely due to wider maximum width of blade at 4.8m for the GW155-4.5MW compared to the 2.8m of the GW165-4.0MW turbine.

Within arrays with both turbines there is slightly more expected annual mortality assigned to the GW155-4.5MW than the GW165-4.0MW, due to the wider maximum blade having a larger percentage risk of collision. Overall the GW165-4.0MW has more estimated annual mortality, this is likely due to there being 135 of these turbines compared to only 20 of the GW155-4.5MW.

**Table 7: Estimated Annual Mortality – 95% Avoidance**

Array	Total Estimated Annual Mortality 95% Avoidance	Estimated Annual Mortality 95% Avoidance GW165 – 4.0MW	Estimated Annual Mortality 95% Avoidance GW155 – 4.5MW
Ban Dakdonna Array	0.069	0.032	0.037
Dak Cheung village Array	0.018	0.018	-
East Central Array	0.116	0.054	0.062
North West Array	0.030	0.030	-
South West Array	0.102	0.047	0.054
Southernmost Array	0.000	0.000	0.000
<b>Total</b>	<b>0.335</b>	<b>0.181</b>	<b>0.154</b>

**Table 8: Estimated Annual Mortality – 0% Avoidance**

Array	Total Estimated Annual Mortality 0% Avoidance	Estimated Annual Mortality 0% Avoidance GW165 – 4.0MW	Estimated Annual Mortality 0% Avoidance GW155 – 4.5MW
Ban Dakdonna Array	1.376	0.640	0.736
Dak Cheung village Array	0.352	0.352	-
East Central Array	2.329	1.0827	1.246
North West Array	0.603	0.603	-
South West Array	2.033	0.945	1.088
Southernmost Array	0	0	0
<b>Total</b>	<b>6.693</b>	<b>3.622</b>	<b>3.071</b>

**Table 9: Collision Risk Percentage turbine comparison**

Collision Risk %	GW165 – 4.0MW	GW155 – 4.5MW
Collision risk factoring in turbine availability (%)	4.02	5.43
Collision risk (%)	4.23	5.72

## 3.2 Potential Biological Removal of Grey-faced buzzard

The results of the calculations for PBR are shown in table 10. The worst case 10,000 individual population assumption indicates a maximum removal from the global population of 210 Grey-faced buzzard per annum before significant population effects occur. The middle case 50,000 gives a PBR of 1750 per annum. The 100,000 estimate allows for non-natural mortality of 3499 individuals per annum.

**Table 10: Potential Biological Removal for different estimated Grey-face buzzard populations**

Population Estimate	100,000 individuals	50,000 individuals	10,000 individuals
$\lambda_{max}$	1.152	1.152	1.152
$R_{max}$	0.152	0.152	0.152
$N_{min}$	91924.74	45962.37	9192.47
Recovery factor	0.5	0.5	0.3
<b>PBR</b>	<b>3499</b>	<b>1750</b>	<b>210</b>

## 4. SUMMARY

### 4.1 Interpreting Estimated Annual Mortality and Potential Biological Removal of Grey-faced buzzard

The results within section 3.1 and 3.2 can be utilised together to show the overall impact the Monsoon Wind Power Project is likely to have on the global population of Grey-faced buzzard. Table 11 shows percentage of PBR estimated to be caused by this project.

In the worst case scenario of 10,000 individuals the project is likely to cause 0.16% of the total annual global non-natural mortalities that could occur before significant negative impacts on the global population occur.

In the middle case 50,000 individuals the project is likely to cause 0.019% of the total annual global non-natural mortalities that could occur before significant negative impacts on the global population occur.

In the 100,000 individuals estimate the project is likely to cause 0.009% of the total annual global non-natural mortalities that could occur before significant negative impacts on the global population occur.

**Table 11: Impacts of Estimated Annual Mortality and Potential Biological Removal**

Population Estimate	100,000 individuals	50,000 individuals	10,000 individuals
Annual Estimated Mortality – 95% avoidance	0.335	0.335	0.335
PBR	3499	1750	210
Percentage of PBR used by Monsoon Wind Project	0.009%	0.019%	0.16%

## 5. REFERENCES

AZ Animals. 2022. *Harris Hawk*. [online] Available at: <<https://a-z-animals.com/animals/harris-hawk/>> [Accessed 14 February 2022].

BirdLife International, 2021. *Butastur indicus*. *The IUCN Red List of Threatened Species 2021: e.T22695726A202433645*. Available at: <<https://dx.doi.org/10.2305/IUCN.UK.2021-3.RLTS.T22695726A202433645.en>> [Accessed 14 February 2022]

BirdLife International, 2022. *Species factsheet: Butastur indicus*. Available at: <<http://datazone.birdlife.org/species/factsheet/22695726>> [Accessed 14 February 2022]

BTO BirdFacts, 2022. *BTO BirdFacts | Buzzard*. [online] Available at: <<https://app.bto.org/birdfacts/results/bob2870.htm>> [Accessed 14 February 2022].

Dillingham, P.W. and Fletcher, D. 2008. Estimating the ability of birds to sustain additional human caused mortalities using a simple decision rule and allometric relationships. *Biological Conservation*. 141: 1783-1792.

Ferguson-Lees, J., and D. A. Christie. 2001. *Raptors of the World*. Christopher Helm, London, UK.

NatureScot. 2000. *Wind farm impacts on birds - Calculating a theoretical collision risk assuming no avoiding action*. [online] Available at: <<https://www.nature.scot/doc/wind-farm-impacts-birds-calculating-theoretical-collision-risk-assuming-no-avoiding-action>> [Accessed 14 February 2022].

Timeanddate.com, 2022. *Sun & moon times today, Pakse, Laos*. [online] Timeanddate.com. Available at: <<https://www.timeanddate.com/astronomy/laos/pakse>> [Accessed 14 February 2022].

Wade, P.R. 1998. Calculating limits to the allowable human-caused mortality of cetaceans and pinnipeds. *Marine Mammal Science* 14: 1–37.

---

**ERM has over 160 offices across the following countries and territories worldwide**

Argentina	The Netherlands
Australia	New Zealand
Belgium	Norway
Brazil	Panama
Canada	Peru
Chile	Poland
China	Portugal
Colombia	Puerto Rico
France	Romania
Germany	Russia
Ghana	Senegal
Guyana	Singapore
Hong Kong	South Africa
India	South Korea
Indonesia	Spain
Ireland	Sweden
Italy	Switzerland
Japan	Taiwan
Kazakhstan	Tanzania
Kenya	Thailand
Malaysia	UAE
Mexico	UK
Mozambique	US
Myanmar	Vietnam

**ERM's London Office**

2<sup>nd</sup> Floor, Exchequer Court  
33 St Mary Axe  
London  
EC3A 8AA

[www.erm.com](http://www.erm.com)

# THE CRM DISCUSSION DOCUMENTS

[Double-click to import picture]



# Monsoon Wind Power Project, Sekong and Attapue Provinces, Lao PDR

## Collision Risk Approach

11 January 2022

Project No.: 0598121

Document details	
Document title	Monsoon Wind Power Project, Sekong and Attapue Provinces, Lao PDR
Document subtitle	Collision Risk Approach
Project No.	0598121
Date	11 January 2022
Version	1.0
Author	Les Hatton
Client Name	Impact Energy Asia Development Limited (IEAD)

#### Document history

Version	Revision	Author	Reviewed by	ERM approval to issue		Comments
				Name	Date	
Preliminary version	00	Les Hatton	Les Hatton	Kamonthip Ma-Oon	11.01.2022	-

---

## Signature Page

11 January 2022

# Monsoon Wind Power Project, Sekong and Attapue Provinces, Lao PDR

## Collision Risk Approach



---

Kamonthip Ma-oon  
Partner

ERM-Siam Co., Ltd.

179 Bangkok City Tower 24th Floor | South Sathorn Road,  
Thungmahamek, Sathorn, Bangkok 10120 | Thailand |

© Copyright 2022 by ERM Worldwide Group Ltd and/or its affiliates ("ERM").  
All rights reserved. No part of this work may be reproduced or transmitted in any form,  
or by any means, without the prior written permission of ERM.

## CONTENTS

<b>1. INTRODUCTION .....</b>	<b>2</b>
1.1 Purpose and scope of this report .....	2
<b>2. METHODOLOGY AND DATA .....</b>	<b>2</b>
2.1 Vantage Point Surveys .....	2
2.1.1 Species Identified.....	3
2.1.2 Approach to high value species .....	4
2.2 Approach to least concern species .....	5
<b>3. CONCLUSIONS AND RECOMMENDATIONS.....</b>	<b>6</b>
3.1 Summary of the Key Findings.....	6
3.2 Conclusion .....	6
<b>4. REFERENCES .....</b>	<b>8</b>

### List of Tables

Table 2.1	Vantage Point relationship to arrays .....	2
Table 2.2	Species recorded during VP surveys.....	4
Table 2.3	Total all flight time in seconds.....	5

### List of Figures

Figure 2.1	Vantage Point and Array Locations .....	3
------------	---	---

## 1. INTRODUCTION

### 1.1 Purpose and scope of this report

This report outlines the proposed approach to assessing the collision risk to birds, based on a review of the reports and datasheets received from Dr. Santi Xayyasith and his bird survey team. The report identifies the species recorded, their conservation status, level of flight activity, migratory or resident status and occurrence patterns. On the basis of this information, it provides a view on whether further assessment using collision risk modelling would be required.

The report provides information on the survey work undertaken and the methodological approach to survey and assessment.

## 2. METHODOLOGY AND DATA


### 2.1 Vantage Point Surveys

Vantage point (VP) surveys were designed according to good international practice, particularly that produced by NatureScot (formerly Scottish Natural Heritage). Given the large and discontinuous area occupied by the arrays, and uncertainties over specific turbine locations, a sampling rather than complete survey approach was adopted. A total of 14 VP's were selected to provide sampling coverage over the habitats associated with the different array areas, including two VP's dedicated to monitoring the flight activity associated with the proposed high voltage (HV) transmission line at the Lao border (VP's 9 and 10).

**Table 2.1 Vantage Point relationship to arrays**

Location	Vantage Points
East Central Arrays	1a, 3a, 2, 4
South East Array	5, 7a, 8
Ban Dakdonna Array	6, 11
Dak Cheung village Array	12
North West Array	13
Southernmost array	14
HV transmission line	9 & 10

Figure 2.1 Vantage Point and Array Locations



Vantage point surveys were planned to have 12 hours survey per VP per month from December 2020 to November 2021 inclusive. Despite the challenges of the terrain, remoteness of location, weather and COVID-19 outbreaks, the survey team completed all the visits to all VP's in every month except April and May, when COVID-19 lockdown restrictions prevented surveys. Total survey time at all VP's was 120 hours, with the exception of VP's 12, 13, and 14 which were added a month after surveys started in response to layout changes, and where survey time was 108 hours.

### 2.1.1 Species Identified

A total of 24 species (excluding three flights to two unspecified species) were recorded. **Table 2.2** lists these in alphabetical order. Of the species recorded, all were raptors, with the exception of two heron species (Chinese Pond Heron and Cinnamon Bittern), a wader (Red-wattled Lapwing) and Greater Hornbill.

All but three of the species recorded were IUCN Least Concern (LC), with two Near Threatened (NT) and one Vulnerable (VU) species recorded.

The majority of species were resident (13), although 11 species were migrants. These proved to be broad front migrants, and there are no Important Bird Areas designated for congregatory or migratory species identified within 50km of the project boundary.

**Table 2.2 Species recorded during VP surveys**

Species common name	Scientific name	IUCN red list status	Resident/Migratory
Besra	<i>Accipiter virgatus</i>	LC	Altitudinal migrant
Black baza	<i>Aviceda leuphotes</i>	LC	Migrant
Black eagle	<i>Ictinaetus malaiensis</i>	LC	Resident
Black-winged kite	<i>Elanus caeruleus</i>	LC	Resident
Changeable hawk-eagle	<i>Nisaetus cirrhatus</i>	LC	Resident
Chinese pond heron	<i>Aredola bacchus</i>	LC	Migrant
Cinnamon bittern	<i>Ixobrychus cinnamomeus</i>	LC	Migrant
Crested goshawk	<i>Accipiter trivirgatus</i>	LC	Resident
Crested serpent eagle	<i>Spilornis cheela</i>	LC	Resident
Eastern buzzard	<i>Buteo japonicas</i>	LC	Migrant
Eurasian kestrel	<i>Falco tinnunculus</i>	LC	Migrant
Eurasian Sparrowhawk	<i>Accipiter nisus</i>	LC	Migrant
Great Hornbill	<i>Buceros bicornis</i>	VU	Resident
Grey-faced Buzzard	<i>Butastur indicus</i>	LC	Migrant
Japanese Sparrowhawk	<i>Accipiter gularis</i>	LC	Migrant
Jerdon's Baza	<i>Aviceda jerdoni</i>	LC	Resident
Mountain Hawk-Eagle	<i>Nisaetus nipalensis</i>	NT	Resident
Northern Goshawk	<i>Accipiter gentilis</i>	LC	Migrant
Oriental Hobby	<i>Falco severus</i>	LC	Resident
Oriental Honey Buzzard	<i>Pernis ptilorhynchus</i>	LC	Resident
Osprey	<i>Pandion haliaetus</i>	LC	Migrant
Red-wattled Lapwing	<i>Vanellus indicus</i>	LC	Resident
Rufous-bellied Eagle	<i>Lophotriorchis kienerii</i>	NT	Resident
Shikra	<i>Accipiter badius</i>	LC	Resident

### 2.1.2 Approach to high value species

Three species with higher IUCN conservation status were identified during the VP surveys. These were the Great Hornbill (VU), the Mountain Hawk-eagle (NT), and the Rufous-bellied Eagle (NT).

The Great Hornbill was observed once from VP10, an area where the high voltage transmission line to Vietnam will be constructed. A total of 15 seconds at collision risk height was recorded. The species is largely sedentary, favouring unlogged evergreen and mixed deciduous woodland. A collision risk model would conclude, on the basis of the observed activity, that no collision would occur. It's presence in an area where HV transmission lines are proposed would indicate a need for additional mitigation, such as flight diverters, to be considered.

Mountain Hawk-eagle was observed for three flights from VP4 and one flight at VP2, both VP's related to the East Central Arrays. Of these four flights, only that at VP2 had flight activity at collision risk height, with a total of 120 seconds recorded. The collision risk modelling would identify the total aerial density based on the coverage of the four VP's sampling the East Central Arrays, divided by the total survey time. This would give an aerial occupancy rate of 4.41925E-07 (i.e. 0.0000041925). Previous experience of collision risk modelling indicates this level of aerial occupancy would not trigger a meaningful collision risk within the lifetime of the wind farm.

Rufous-bellied eagle was also recorded once at VP4 and once at VP2. Of these two flights 30 seconds was spent at collision risk height at VP2, and none at VP4. Likely collision risk is therefore, as with mountain hawk-eagle, likely to be statistically insignificant.

## 2.2 Approach to least concern species

All other species recorded were least concern. Table 2.3 indicates the total flight time, at all heights, recorded between December 2020 and November 2021 inclusive. The proportion of flights at collision risk height is much smaller.

**Table 2.3 Total all flight time in seconds**

Species common name	Altitudinal migrant	Migrant	Resident	Time at CRH
Besra	415			270
Black Baza		120		30
Black Eagle			8262	3450
Black-winged kite			600	300
Changeable Hawk-eagle			72	0
Chinese Pond Heron		90		0
Cinamon Bittern		120		0
Crested Goshawk			1506	465
Crested Serpent Eagle			5105	975
Eastern Buzzard		120		120
Eurasian Kestrel		610		90
Eurasian Sparrowhawk		150		15
Great Hornbill (VU)			135	15
Grey-faced Buzzard		3829		2190
Japanese Sparrowhawk		90		45
Jerdon's Baza			915	450
Mountain Hawk-Eagle (NT)			300	120
Northern Goshawk		82		45
Oriental Hobby			924	375
Oriental Honey Buzzard			2310	1110
Osprey		600		375
Red-wattled Lapwing			30	0
Rufous-bellied Eagle (NT)			180	60
Shikra			1288	180

Of the LC species, no flights at collision risk height were recorded for Chinese Pond Heron, Cinnamon Bittern, Changeable Hawk-eagle, and Red-wattled Lapwing. These have therefore been ruled out from further assessment.

Most migrant species spent less than 375 seconds at collision risk height, and this would not generate sufficient time at collision risk height to have a statistical probability of a collision within the lifetime of the wind farm. Grey-faced Buzzard which was commonly encountered at ten of the fourteen VP's from November through till March, was observed for a total of 2190 seconds at collision risk height. Observation time for the five migration months was 600 hours. Over all the arrays that it occurs, the likelihood of collision is still small, as the total density is likely to be less than 0.0000612

hectare/hours. The statistical probability of a collision is therefore likely to be very low. The population is not known, but is assumed to exceed 100,000 individuals (Birdlife International 2021), and it has a large area of occupancy. It is therefore highly unlikely that the wind farm would have any population level effect.

Generally, resident birds, by virtue of being present throughout the year, have a larger number of observations. Most are below a thousand seconds at collision risk height, and therefore unlikely to trigger a likely significant collision risk within the life time of the wind farm, and all are from widespread and common populations. Three species have elevated times at collision risk time, most notably Black Eagle (3450 seconds in 1320 hours of observation), Crested Serpent Eagle (975 seconds in 1416 hours of observation), and Oriental Honey Buzzard (1110 seconds in 960 hours of observation).

Black Eagle is widespread within the wind farm, having been recorded at eleven VP's. Actual aerial occupancy is still relatively low (approximately 0.0000127 hectare/hours), however, over all the arrays there is potential for collisions to result in some local reduction in breeding pairs, but this is unlikely to be significant beyond this local scale, given the wide distribution of the species, and a population believed to be in excess of 10,000 individuals within its extensive range.

Crested Serpent Eagle is even more widespread, having been observed at twelve VP locations. Activity at collision risk height is lower, and therefore the overall likelihood of collision is lower. Some arrays, most notably the East Central Arrays, have higher levels of activity, and the possibility of local effects cannot be excluded. However it is a widespread and common species over much of its range, with a stable population.

Similarly the Oriental honey buzzard is also widespread, being recorded at ten VP's, and has activity at collision risk height very similar to the Crested Serpent Eagle. Risks to this population are slight, as it has an even bigger global range than the two previous species, and may have a population of 100,000-1,000,000 individuals, although the trend is believed to be declining.

### 3. CONCLUSIONS AND RECOMMENDATIONS

#### 3.1 Summary of the Key Findings

Based on a high level of sampling effort over the different geographical arrays within the proposed Monsoon wind farm there is no significant risk to any higher risk species (NT or VU) recorded during the VP surveys.

All migrant species are IUCN LC, and most record levels of flight at collision risk height that are unlikely to result in a collision risk model indicating a risk of collision within the life time of the wind farm. This is primarily due to the low proportion of collision risk flights in comparison to the survey area and hours of observation. The only migrant with relatively high levels of flight at collision risk height is the Grey-faced Buzzard, and as this still amounts to only 2190 seconds over 1188 hours of observation, the risk of collision remains low. This together with the large and widespread population indicates no population level effect is likely.

Resident birds tend to have higher levels of recorded flight activity, but of these, only three came close to activity levels around or greater than 1000 seconds. None of these three species were likely to exceed anything other than local effects, due to the low statistical likelihood of collision, and widespread and common nature of the populations.

#### 3.2 Conclusion

No higher value species (NT, VU) occur at levels likely to trigger collision risk, and aerial occupancy rates are so low that there is little to no value in running a collision risk model.

This is also largely true for LC species, although the possibility of collisions for species such as Black Eagle cannot be ruled out. However any such effects would only ever be significant, if at all, at the

local population level and no significant population level effect is anticipated, given the range and population status of the LC species.

There is some evidence of broad front raptor migration, but all species involved are LC, and again the low levels of likely aerial occupancy indicate a collision risk model would be unlikely to indicate sufficient collisions within the lifetime of the wind farm to threaten the status of such populations. No Important Bird Areas for migration are found within 50km of the proposed wind farm and extensive field observations during the important migration months do not indicate the study area is a significant migratory or congregatory area.

No cumulative impact assessment has been performed, however there are no other wind farms in the Lao area and the spatial separation from transboundary wind farms e.g. in Vietnam, would suggest little, if any, likelihood of cumulative effects with other wind farms.

On this basis it is not proposed to undertake further collision risk modelling, although the data to do so is available should it be required.

## 4. REFERENCES

BirdLife International. 2016a. *Ictinaetus malaiensis*. *The IUCN Red List of Threatened Species* 2016: e.T22696019A93538909. <https://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T22696019A93538909.en>. Accessed on 10 January 2022.

BirdLife International. 2016b. *Spilornis cheela*. *The IUCN Red List of Threatened Species* 2016: e.T22695293A95221642. <https://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T22695293A95221642.en>. Accessed on 10 January 2022.

BirdLife International. 2021a. *Butastur indicus*. *The IUCN Red List of Threatened Species* 2021: e.T22695726A202433645. <https://dx.doi.org/10.2305/IUCN.UK.2021-3.RLTS.T22695726A202433645.en>. Accessed on 10 January 2022.

BirdLife International. 2021b. *Pernis ptilorhynchus*. *The IUCN Red List of Threatened Species* 2021: e.T22694995A199637824. <https://dx.doi.org/10.2305/IUCN.UK.2021-3.RLTS.T22694995A199637824.en>. Accessed on 10 January 2022.

Scottish Natural Heritage. March 2017. Recommended bird survey methods to inform impact assessment of onshore wind farms.

---

**ERM has over 160 offices across the following countries and territories worldwide**

Argentina	The Netherlands
Australia	New Zealand
Belgium	Norway
Brazil	Panama
Canada	Peru
Chile	Poland
China	Portugal
Colombia	Puerto Rico
France	Romania
Germany	Russia
Ghana	Senegal
Guyana	Singapore
Hong Kong	South Africa
India	South Korea
Indonesia	Spain
Ireland	Sweden
Italy	Switzerland
Japan	Taiwan
Kazakhstan	Tanzania
Kenya	Thailand
Malaysia	UAE
Mexico	UK
Mozambique	US
Myanmar	Vietnam

**ERM-Siam Co., Ltd.**

179 Bangkok City Tower 24th Floor,  
South Sathorn Road, Thungmahamek,  
Sathorn, Bangkok 10120, Thailand

T: (662) 679 5200

[www.erm.com](http://www.erm.com)